

Emergency Department		
Description	ICD10-CM	Documentation Tips
<b>ABDOMINAL PAIN</b>		
	<b>Code</b>	<b>Tip</b>
Left upper quadrant pain	R10.12	Indicate pain, tenderness or rebound tenderness
Right lower quadrant pain	R10.31	
Upper abdominal pain, unspecified	R10.10	
Right lower quadrant rebound abdominal tenderness	R10.823	Indicate location: quadrant, epigastric, umbilical, generalized
<b>ATRIAL FIBRILLATION</b>		
	<b>Code</b>	<b>Tip</b>
Paroxysmal Afib	I48.0	Indicate fibrillation vs. flutter, paroxysmal, persistent, chronic, typical or atypical
Chronic Afib	I48.2	
Typical Atrial Flutter	I48.3	
Atypical Atrial Flutter	I48.4	
<b>ASTHMA</b>		
	<b>Code</b>	<b>Tip</b>
Mild intermittent	J45.20	Indicate mild, moderate, severe intermittent, persistent status asthmaticus acute exacerbation
Mod persistent w/status asthmaticus	J45.42	
Severe persistent w/acute exacerbation	J45.51	
Persistent Afib	I48.1	
<b>CHF</b>		
	<b>Code</b>	<b>Tip</b>
Acute Systolic CHF	I50.21	Indicate systolic, diastolic, or combined and acute, chronic or acute on chronic
Chronic Diastolic CHF	I50.32	
<b>COPD</b>		
	<b>Code</b>	<b>Tip</b>
COPD w/acute lower resp. infection	J44.0	Indicate w/acute lower respiratory infection or w/acute exacerbation
COPD with acute exacerbation	J44.1	
Chronic obstructive pulmonary	J44.9	

Emergency Department		
Description	ICD10-CM	Documentation Tips
<b>DIABETES</b>		
	<b>Code</b>	<b>Tip</b>
Type I No complications	E11.9	Indicate Type I or Type II and all manifestations. Use linking statement (i.e. diabetic cataract)
Type 2 diabetes mellitus with hyperosmolarity with coma	E11.01	
<b>FRACTURE</b>		
	<b>Code</b>	<b>Tip</b>
Stress fracture, hip, unspecified, initial encounter for fracture	M84.359A	Indicate right or left, specific area of bone, type of fracture, severity of fracture, episode of care
Non displaced fracture of coracoid process, left shoulder, subsequent encounter for fracture with delayed healing	S42.135G	
<b>PNEUMONIA</b>		
	<b>Code</b>	<b>Tip</b>
RSV Pneumonia	J12.1	Indicate viral or bacterial and specific causal organism
Adenoviral Pneumonia	J12.0	
Pneumonia due to MSSA	J15.211	
<b>RESPIRATORY FAILURE</b>		
	<b>Code</b>	<b>Tip</b>
Acute RF with hypoxia	J96.01	Indicate acute, chronic or acute on chronic. Also indicate hypercapnia or hypoxia. If resting or expected to be on vent, or on vent for airway protection, do not use the term Respiratory Failure
Acute RF with hypercapnia	J96.02	
Chronic RF with hypoxia	J96.11	
<b>SEIZURES</b>		
	<b>Code</b>	<b>Tip</b>
Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes w/ seizures of localized onset, not intractable, w/ status epilepticus	G40.001	Indicate recurrent or single episode; intractable or not-intractable; localized or generalized, with or without status epilepticus
Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes w/ seizures of localized onset, intractable, w/o status epilepticus	G40.019	
Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, not intractable, w/o status epilepticus	G40.209	
Juvenile myoclonic epilepsy, not intractable, with status epilepticus	G40.B01	

\* Indicate Episode of Care (initial, subsequent, sequela) for injuries or fractures  
 \* Always indicate left, right or bilateral  
**These are intended to be examples not a comprehensive listing. Please refer to the official ICD-10-CM code set for details**

October 2013-September 2015 except limited code updates for new technology and new diseases

Questions: [BHICD10ProviderQA@bannerhealth.com](mailto:BHICD10ProviderQA@bannerhealth.com)