
 OB GYN		
Description	ICD10-CM	Documentation Tips
Abdominal/Pelvic Pain	Code	Tip
Left upper quadrant pain	R10.12	Indicate pain, tenderness or rebound tenderness Indicate location: quadrant, epigastric, umbilical, generalized
Right lower quadrant pain	R10.31	
Right lower quadrant rebound abdominal tenderness	R10.823	
Pelvic Pain	R10.2	
Cervical Dysplasia	Code	Tip
Mild cervical dysplasia	N87.0	Indicate CIN I, CIN II, CIN III (mild, moderate, severe)
Moderate cervical dysplasia	N87.1	
Dysplasia of cervix uteri, unspecified	N87.9	
Depression	Code	Tip
Puerperal psychosis	F53	Indicate if Post Partem; indicate mild, moderate, severe; single episode or recurrent
Major depressive disorder, single	F32.1	
Major depressive disorder, recurrent, in full remission	F33.42	
Diabetes	Code	Tip
Type I No complications	E10.9	Indicate Type I or Type II and all manifestations. Use linking statement; Gestational=indicate diet controlled or uncontrolled, insulin
TYPE II No complication	E11.9	
Gestational diabetes mellitus in	O24.410	
Hypertension	Code	Tip
Essential (primary) hypertension	I10	Gestational or pre-existing; level of severity; gestational age
Pre-existing hypertension with pre-eclampsia, first trimester	O11.1	
Gestational (pregnancy-induced) hypertension w/o significant	O13.3	

<div> OB GYN</div>		
Description	ICD10-CM	Documentation Tips
Multiple Gestations	Code	Tip
Twin pregnancy, monochorionic/monoamniotic, first trimester	O30.011	Indicate number of fetuses; Number of placentas and number of amniotic sacs; gestational age
Triplet pregnancy with two or more monoamniotic fetuses, second trimester	O30.122	
Quadruplet pregnancy, unable to determine number of placenta and number of amniotic sacs, third trimester	O30.293	
Premature ROM	Code	Tip
Preterm premature ROMs, onset of labor within 24 hours of rupture, third trimester	O42.013	Indicate time between rupture and onset of labor; gestational age
Full-term premature ROMs, onset of labor within 24 hours of rupture	O42.02	
Pregnancy	Code	Tip
Encounter for supervision of normal first pregnancy, first trimester	Z34.01	Indicate gestational age at every visit for episode of care (1st, 2nd, 3rd trimester), childbirth or postpartum Use on maternal record only
Supervision of pregnancy with history of pre-term labor, third trimester	O09.213	
Placenta previa with hemorrhage, third trimester	O44.13	
Uterine Prolapse	Code	Tip
Incomplete uterovaginal prolapse	N81.2	Indicate degree (1st, 2nd, 3rd, or complete)
Complete uterovaginal prolapse	N81.3	
Maternal care for prolapse of gravid uterus, second trimester	O34.522	
UTI	Code	Tip
Acute cystitis with hematuria	N30.01	Indicate location of UTI [ureter, bladder, urethra]. For bladder indicate acute, interstitial, chronic, trigonitis, irradiation and with or without hematuria
Chronic cystitis without hematuria	N30.20	
Urethritis	N34.2	
Weight/Obesity	Code	Tip
Excessive weight gain in pregnancy, 3rd trimester	O26.03	Indicate overweight, obese, or morbidly obese and BMI
Low weight gain in pregnancy, 2nd trimester	O26.12	
Obesity complicating childbirth	O99.214	
<div><div>* Indicate Episode of Care (initial, subsequent, sequela) for injuries or fractures</div><div>* Always indicate left, right or bilateral</div><div>* These are intended to be examples not a comprehensive listing. Please refer to the official ICD-10-CM code set for details</div></div>		
<div>October 2013-September 2015 except limited code updates for new technology and new diseases</div> <div>Questions: BHICD10ProviderQA@bannerhealth.com</div>		