
 Genitourinary		
Description	ICD10-CM	Documentation Tips
BPH	Code	Tip
Enlarged prostate with lower	N40.1	Enlarged Prostate vs. Nodular Prostate; with or without LUTS Use additional code for associated symptoms when specified, such as,
Nodular prostate without lower	N40.2	
Nodular prostate with lower urinary tract symptoms		
ED	Code	Tip
Erectile dysfunction due to arterial insufficiency	N52.01	Indicate due to, such as, post surgery (type of surgery) or due to underlying disease
Drug-induced erectile dysfunction	N52.2	
Hepatitis	Code	Tip
Hepatitis A without hepatic coma	B15.9	Indicate acute or chronic; Cause (viral, or alcoholic); Type Indicate with or without hepatic coma
Acute hepatitis B with delta-agent with hepatic coma	B16.0	
Acute hepatitis C without hepatic coma	B17.10	
Kidney Failure	Code	Tip
AKF with tub necrosis	N17.0	Indicate acute with tubular, cortical, or medullary necrosis. If chronic, indicate stage
AKF with med necrosis	N17.2	
CKD stage 3	N18.3	
ESRD	N18.6	

 Genitourinary		
Description	ICD10-CM	Documentation Tips
Pain, Female Genitourinary	Code	Tip
Dyspareunia	N94.1	
Primary dysmenorrhea	N94.4	
Sepsis	Code	Tip
Unspecified Organism	A41.9	Indicate causal organism, severe sepsis, septic shock & organ failure. "Urosepsis" is not specific & should be avoided
Sepsis due to E.coli	A41.51	
Severe Sepsis	R65.20	
UTI	Code	Tip
Acute cystitis with hematuria	N30.01	Indicate location of UTI [ureter, bladder, urethra]. For bladder indicate acute, interstitial, chronic, trigonitis, irradiation and with or without hematuria
Chronic cystitis without hematuria	N30.20	

* Always indicate left, right or bilateral
* <u>These are intended to be examples not a comprehensive listing. Please refer to the official ICD-10-CM code set for details</u>
October 2013-September 2015 except limited code updates for new technology and new diseases
Questions: BHICD10ProviderQA@bannerhealth.com