

RAF HCC			
Description	ICD10-CM	HCC	Documentation Tips
Diabetes Type 1	Code	Model V22	Tip
Type 1 No complications	E10.9	19	Indicate Type 1 or Type 2 and all manifestations. Use linking statement (i.e. diabetic cataract); with or w/o coma, due to
Type 1 diabetes mellitus with hyperglycemia	E10.65	18	
Type 1 diabetes mellitus with diabetic chronic kidney disease	E10.22	18	
Diabetes Type 2	Code	Model V22	Tip
Type 2 No complication	E11.9	19	Indicate Type 1 or Type 2 and all manifestations. Use linking statement (i.e. diabetic cataract); with or w/o coma, due to Report long term (current) use of insulin Z79.4
Type 2 diabetes mellitus with hyperglycemia	E11.65	18	
Type 2 diabetes mellitus with diabetic nephropathy	E11.21	18	
Type 2 diabetes mellitus with unspecified diabetic retinopathy with macular edema	E11.311	18	
Type 2 diabetes mellitus with proliferative diabetic retinopathy without macular edema	E11.359	18, 122	
Type 2 diabetes mellitus with diabetic polyneuropathy	E11.42	18	
Type 2 diabetes mellitus with diabetic peripheral angiopathy without gangrene	E11.51	18, 108	
COPD	Code	Model V22	Tip
COPD w/acute lower resp.	J44.0	111	Indicate w/acute lower respiratory infection or w/acute exacerbation
COPD w/ acute exacerbation	J44.1	111	
Chronic obstructive pulmonary disease, unspecified	J44.9	111	
Respiratory Failure	Code	Model V22	Tip
Acute RF with hypoxia	J96.01	84	Indicate acute, chronic or acute on chronic. Also indicate hypercapnia or hypoxia. If resting or expected to be on vent, or on vent
Acute RF with hypercapnia	J96.02	84	
Chronic RF with hypoxia	J96.11	84	
Obesity	Code	Model V22	Tip
Morbid Obesity due to excess calories	E66.01	22	Indicate morbid, drug induced, or overweight. Document BMI
(BMI) 40.0-44.9, adult	Z68.41	22	
(BMI) 45.0-49.9, adult	Z68.42	22	
(BMI) 50-59.9, adult	Z68.43	22	
(BMI) 60.0-69.9, adult	Z68.44	22	
(BMI) 70 or greater, adult	Z68.45	22	

These are intended to be examples not a comprehensive listing. Please refer to the official ICD-10-CM code set for additional diagnosis.

RAF HCC

Description	ICD10-CM	HCC	Documentation Tips
Breast Cancer	Code	Model V22	Tip
Malignant neoplasm of nipple and areola, right female breast	C50.011	12	Indicate primary, secondary, or CA in situ; specific location
Malignant neoplasm of lower-outer quadrant of left female breast	C50.512	12	
Secondary malignant neoplasm of breast	C79.81	10	
Prostate Cancer	Code	Model V22	Tip
Malignant neoplasm of prostate	C61	12	Indicate primary, secondary, or CA in situ
Secondary malignant neoplasm of genital organs	C79.82	10	
Lymphoma	Code	Model V22	Tip
Anaplastic large cell lymphoma, ALK-negative, intra-abdominal lymph nodes	C84.73	10	Indicate specific type (Hodgkin, follicular, non-Hodgkin); location (head, neck, axilla, inguinal, intrapelvic, spleen, etc)
Lymphocyte-rich classical Hodgkin lymphoma, spleen	C81.47	10	
Cutaneous T-cell lymphoma, unspecified lymph nodes of head, face, and neck	C84.A1	10	
Leukemia	Code	Model V22	Tip
Chronic myeloid leukemia, BCR/ABL-positive, in remission	C92.11	9	Indicate type (lymphoid, myeloid, monocytic); in remission, or in relapse
Acute myeloblastic leukemia, not having achieved remission	C92.00	8	
Ulcer	Code	Model V22	Tip
Pressure ulcer of right elbow, unstageable	L89.010	158	Pressure vs. non-pressure; stage; location and laterality
Pressure ulcer of right upper back, stage 3	L89.113	158	
Non-pressure chronic ulcer of left calf limited to breakdown of skin	L97.221	161	
Non-pressure chronic ulcer of right ankle with necrosis of bone	L97.314	161	
Non-pressure chronic ulcer of buttock with fat layer exposed	L97.412	161	

These are intended to be examples not a comprehensive listing. Please refer to the official ICD-10-CM code set for additional diagnosis.

RAF HCC

Description	ICD10-CM	HCC	Documentation Tips
Alcohol/Drug Use, Abuse or Dependence	Code	Model V22	Tip
Alcohol abuse with alcohol-induced psychotic disorder with delusions	F10.150	54	Indicate use, abuse, or dependence; with intoxication, anxiety disorder, in remission, or withdrawal; uncomplicated; indicate specific drug; Evidence of alcohol involvement determined by blood alcohol level (BAL): include level if known
Alcohol dependence, in remission	F10.21	55	
Alcohol dependence with withdrawal, unspecified	F10.239	55	
Bipolar	Code	Model V22	Tip
Bipolar disorder, current episode hypomanic	F31.0	58	Indicate if current or partial episode or, in full remission; mild, moderate or severe
Bipolar disorder, current episode	F31.63	58	
Bipolar disorder, in full remission, most recent episode depressed	F31.76	58	
Depression	Code	Model V22	Tip
Major depressive disorder, single episode, moderate	F32.1	58	Indicate single or recurrent episode; mild, moderate, severe; with or without psychotic features; full or partial remission
Major depressive disorder, single episode, in partial remission	F32.4	58	
Major depressive disorder, recurrent, severe with psychotic symptoms	F33.2	58	
Major depressive disorder, recurrent, severe with psychotic symptoms	F33.3	58	
Major depressive disorder, recurrent, in full remission	F33.42	58	
Atrial Fibrillation	Code	Model V22	Tip
Paroxysmal Afib	I48.0	96	Indicate fibrillation vs. flutter, paroxysmal, persistent, chronic, typical or atypical
Persistent Afib	I48.1	96	
Chronic Afib	I48.2	96	
Typical Atrial Flutter	I48.3	96	
CHF	Code	Model V22	Tip
Acute Systolic CHF	I50.21	85	Indicate systolic, diastolic, or combined and acute, chronic or acute on chronic
Chronic Diastolic CHF	50.32	85	
Acute on chronic combined CHF	I50.43	85	
Chronic Systolic CHF	I50.22	85	
Acute Diastolic CHF	I50.31	85	

These are intended to be examples not a comprehensive listing. Please refer to the official ICD-10-CM code set for additional diagnosis.